

COVID-19 Vaccine Recommendations

Healthy Adults: Administration of Primary Series and Booster Doses (ages 18 and older)

Date: July 19, 2022



	Dose 1		Dose 2		Dose 3		Dose 4
Pfizer Cap: Gray or Purple Ages 18 and older <i>See Child and Adolescent charts for recommendations for those under 18 years of age</i>	(primary) 0.3 mL	Wait 3 – 8 weeks →	(primary) 0.3 mL	Wait 5 months →	(booster 1) Pfizer: 0.3 mL OR Moderna: 0.25 mL OR J&J: 0.5 mL	Wait 4 months → Eligible Healthy Patients: - Ages 50 years and older	(booster 2) Pfizer: 0.3 mL OR Moderna: 0.25 mL
Moderna Cap: Red Ages 18 and older <i>See Child and Adolescent charts for recommendations for those under 18 years of age</i>	(primary) 0.5 mL	Wait 4 – 8 weeks →	(primary) 0.5 mL	Wait 5 months →	(booster 1) Pfizer: 0.3 mL OR Moderna: 0.25 mL OR J&J: 0.5 mL	Wait 4 months → Eligible Healthy Patients: - Ages 50 years and older	(booster 2) Pfizer: 0.3 mL OR Moderna: 0.25 mL
Novavax Ages 18 and older	(primary) 0.5 mL	Wait 3 – 8 weeks →	(primary) 0.5 mL		NOT CURRENTLY RECOMMENDED		NOT CURRENTLY RECOMMENDED
J & J Ages 18 and older	(primary) 0.5 mL		Wait 2 months →		(booster 1) Pfizer: 0.3 mL OR Moderna: 0.25 mL OR J&J: 0.5 mL	Wait 4 months → Eligible Healthy Patients: - Ages 50 years and older - Ages 18–49 years who received 2 previous doses of J&J	(booster 2) Pfizer: 0.3 mL OR Moderna: 0.25 mL NOTE: J&J may NOT be used as a 2nd booster

NOTE: J&J should only be used in limited situations and cannot be used as a second booster.

Please see CDC Interim Clinical Considerations for official guidance.