

COVID-19 Vaccine Recommendations

Administration of Primary Series and Booster Doses for **Immunocompromised** Adults and Adolescents

Date: May 20, 2022



	Dose 1		Dose 2		Additional Dose		Booster 1		Booster 2
Pfizer (gray or purple cap) Ages 12 and older	0.3 mL	Wait 3 – 8 weeks →	0.3 mL	Wait 4 weeks →	Pfizer: 0.3 mL OR Moderna: 0.5 mL	Wait 3 months →	Pfizer: 0.3 mL OR Moderna: 0.25 mL OR J&J: 0.5 mL	Wait 4 months → Eligible Patients: - Everyone ages 50 years and older - Everyone ages 12 years and older who are moderately to severely immunocompromised	Pfizer: 0.3 mL OR Moderna: 0.25 mL
Moderna (red cap) Ages 18 and older	0.5 mL	Wait 4 – 8 weeks →	0.5 mL	Wait 4 weeks →	Pfizer: 0.3 mL OR Moderna: 0.5 mL	Wait 3 months →	Pfizer: 0.3 mL OR Moderna: 0.25 mL OR J&J: 0.5 mL	Wait 4 months → Eligible Patients: - Everyone ages 50 years and older - Everyone ages 18 years and older who are moderately to severely immunocompromised	Pfizer: 0.3 mL OR Moderna: 0.25 mL
J & J Ages 18 and older	0.5 mL	Wait 4 weeks →			Pfizer: 0.3 mL OR Moderna: 0.5 mL	Wait 2 months →	Pfizer: 0.3 mL OR Moderna: 0.25 mL OR J&J: 0.5 mL	Wait 4 months → Eligible Patients: - Everyone ages 50 years and older - Everyone ages 18 years and older who are moderately to severely immunocompromised - Everyone ages 18–49 years who received 2 previous doses of J&J	Pfizer: 0.3 mL OR Moderna: 0.25 mL

NOTE: Pfizer and Moderna are preferred over J&J for primary series and booster doses.