

COVID-19 Vaccine Recommendations

Healthy Adults: Administration of Primary Series and Booster Doses (ages 18 and older)

Date: December 9, 2022



PRIMARY SERIES (monovalent products only)				BOOSTER DOSES (bivalent products only)	
	Dose 1		Dose 2	Booster	
Pfizer Cap: Gray or Purple	(primary) Pfizer Monovalent: 0.3 mL	Wait 3 – 8 weeks →	(primary) Pfizer Monovalent: 0.3 mL	Wait 2 months → (Regardless of previous monovalent booster doses given)	Pfizer Bivalent: 0.3 mL OR Moderna Bivalent: 0.5 mL
Moderna Cap: Red	(primary) Moderna Monovalent: 0.5 mL	Wait 4 – 8 weeks →	(primary) Moderna Monovalent: 0.5 mL	Wait 2 months → (Regardless of previous monovalent booster doses given)	Pfizer Bivalent: 0.3 mL OR Moderna Bivalent: 0.5 mL
Novavax	(primary) Novavax Monovalent: 0.5 mL	Wait 3 – 8 weeks →	(primary) Novavax Monovalent: 0.5 mL	Wait 2 months → (Regardless of previous monovalent booster doses given)	Pfizer Bivalent: 0.3 mL OR Moderna Bivalent: 0.5 mL
J & J J&J should only be used in limited situations	(primary) J&J Monovalent: 0.5 mL			Wait 2 months → (Regardless of previous monovalent booster doses given)	Pfizer Bivalent: 0.3 mL OR Moderna Bivalent: 0.5 mL

See Child and Adolescent charts for recommendations for those under 18 years of age

Please see CDC Interim Clinical Considerations for official guidance.