

# COVID-19 Vaccine Recommendations

**Healthy Adults:** Administration of Primary Series and Booster Doses (ages 18 and older)

Date: October 12, 2022



| PRIMARY SERIES (monovalent products only)                         |  |                          |  | BOOSTER DOSES (bivalent products only)   |   |
|---|--|--------------------------|--|--|---|
|   | Dose 1   |                          | Dose 2   | Booster  |   |
| <b>Pfizer</b><br>Cap: Gray or Purple                              | (primary)<br><b>Pfizer Monovalent:</b><br><b>0.3 mL</b>  | Wait<br>3 – 8 weeks<br>→ | (primary)<br><b>Pfizer Monovalent:</b><br><b>0.3 mL</b>  | Wait<br>2 months<br>→<br>(Regardless of previous monovalent booster doses given) | <b>Pfizer Bivalent: 0.3 mL</b><br>OR<br><b>Moderna Bivalent: 0.5 mL</b> |
| <b>Moderna</b><br>Cap: Red  | (primary)<br><b>Moderna Monovalent:</b><br><b>0.5 mL</b> | Wait<br>4 – 8 weeks<br>→ | (primary)<br><b>Moderna Monovalent:</b><br><b>0.5 mL</b> | Wait<br>2 months<br>→<br>(Regardless of previous monovalent booster doses given) | <b>Pfizer Bivalent: 0.3 mL</b><br>OR<br><b>Moderna Bivalent: 0.5 mL</b> |
| <b>Novavax</b>  | (primary)<br><b>Novavax Monovalent:</b><br><b>0.5 mL</b> | Wait<br>3 – 8 weeks<br>→ | (primary)<br><b>Novavax Monovalent:</b><br><b>0.5 mL</b> | Wait<br>2 months<br>→<br>(Regardless of previous monovalent booster doses given) | <b>Pfizer Bivalent: 0.3 mL</b><br>OR<br><b>Moderna Bivalent: 0.5 mL</b> |
| <b>J &amp; J</b><br>J&J should only be used in limited situations | (primary)<br><b>J&amp;J Monovalent:</b><br><b>0.5 mL</b> |                          |  | Wait<br>2 months<br>→<br>(Regardless of previous monovalent booster doses given) | <b>Pfizer Bivalent: 0.3 mL</b><br>OR<br><b>Moderna Bivalent: 0.5 mL</b> |

See Child and Adolescent charts for recommendations for those under 18 years of age

Please see CDC Interim Clinical Considerations for official guidance.