

COVID-19 Vaccine Recommendations

Immunocompromised Children and Adolescents: Ages 5 years through 11 years

Last Updated: December 20, 2023



Ages: 5 years thru 11 years								
	COVID-19 Vaccine History		2023 – 2024 Updated Vaccines					
Pfizer Cap: Blue Age: 5 – 11 years	Unvaccinated		Pfizer: 0.3 mL	Wait 3 weeks	Pfizer: 0.3 mL	Wait 4 weeks	Pfizer: 0.3 mL	
		OR						
Moderna Cap: Dark Blue Label: Green Age: 6 mo – 11 years			Moderna: 0.25 mL	Wait 4 weeks	Moderna: 0.25 mL	Wait 4 weeks	Moderna: 0.25 mL	
								<i>Additional doses may be administered at least 2 months after last dose of mRNA vaccine.</i>
	1 dose any Pfizer	Wait 3 weeks	Pfizer: 0.3 mL	Wait 4 weeks	Pfizer: 0.3 mL	<i>Additional doses may be administered at least 2 months after last dose of mRNA vaccine.</i>		
	2 doses any Pfizer	Wait 4 weeks	Pfizer: 0.3 mL	<i>Additional doses may be administered at least 2 months after last dose of mRNA vaccine.</i>				
	3 or more doses any mRNA	Wait 8 weeks	Pfizer: 0.3 mL OR Moderna: 0.25 mL	<i>Additional doses may be administered at least 2 months after last dose of mRNA vaccine.</i>				
	1 dose any Moderna	Wait 4 weeks	Moderna: 0.25 mL	Wait 4 weeks	Moderna: 0.25 mL	<i>Additional doses may be administered at least 2 months after last dose of mRNA vaccine.</i>		
2 doses any Moderna	Wait 4 weeks	Moderna: 0.25 mL	<i>Additional doses may be administered at least 2 months after last dose of mRNA vaccine.</i>					
3 or more doses any mRNA	Wait 8 weeks	Pfizer: 0.3 mL OR Moderna: 0.25 mL	<i>Additional doses may be administered at least 2 months after last dose of mRNA vaccine.</i>					

*Heterologous mRNA products may be used for updated vaccines for this age group, **except when completing the first 3 doses (see exceptions below)**.

Mixing and matching Pfizer and Moderna is authorized after dose 3.

Pfizer Additional Doses:

- Pfizer Blue Cap (0.3 mL)

Moderna Additional Doses:

- Moderna Blue Cap/Green Label (0.25 mL)

Interchangeability:

COVID-19 vaccine doses from the same manufacturer should be administered whenever recommended. In the following circumstances, an age-appropriate COVID-19 vaccine from a different manufacturer may be administered:

- Same vaccine not available at the vaccination site at the time of the clinic visit
- Previous dose unknown
- Person would otherwise not receive a recommended vaccine dose
- Person starts but unable to complete a vaccination series with the same COVID-19 vaccine due to a contraindication