

COVID-19 Vaccine Recommendations

Immunocompromised Children and Adolescents: Ages 6 months through 4 years

Last Updated: December 20, 2023



Ages: 6 months thru 4 years								
	COVID-19 Vaccine History		2023 – 2024 Updated Vaccines					
Pfizer Cap: Yellow Age: 6 mo – 4 years	Unvaccinated		Pfizer: 0.3 mL	Wait 3 weeks	Pfizer: 0.3 mL	Wait 8 weeks	Pfizer: 0.3 mL	
		OR						
			Moderna: 0.25 mL	Wait 4 weeks	Moderna: 0.25 mL	Wait 4 weeks	Moderna: 0.25 mL	
Moderna Cap: Dark Blue Label: Green Age: 6 mo – 11 years							Additional doses may be administered at least 2 months after last dose of mRNA vaccine.	
	1 dose any Pfizer	Wait 3 weeks	Pfizer: 0.3 mL	Wait 8 weeks	Pfizer: 0.3 mL	Additional doses may be administered at least 2 months after last dose of mRNA vaccine.		
	2 doses any Pfizer	Wait 8 weeks	Pfizer: 0.3 mL	Additional doses may be administered at least 2 months after last dose of mRNA vaccine.				
	3 or more doses any Pfizer	Wait 8 weeks	Pfizer: 0.3 mL	Additional doses may be administered at least 2 months after last dose of mRNA vaccine.				
	1 dose any Moderna	Wait 4 weeks	Moderna: 0.25 mL	Wait 4 weeks	Moderna: 0.25 mL	Additional doses may be administered at least 2 months after last dose of mRNA vaccine.		
	2 doses any Moderna	Wait 4 weeks	Moderna: 0.25 mL	Additional doses may be administered at least 2 months after last dose of mRNA vaccine.				
	3 doses any Moderna	Wait 8 weeks	Moderna: 0.25 mL	Additional doses may be administered at least 2 months after last dose of mRNA vaccine.				

*Heterologous mRNA products **may NOT** be used for updated vaccines for this age group. Mixing and matching Pfizer and Moderna **is NOT** authorized (except in certain situations, listed below).

Pfizer Additional Doses:

- Pfizer Yellow Cap (0.3 mL)

Moderna Additional Doses:

- Moderna Blue Cap/Green Label (0.25 mL)

Interchangeability:

COVID-19 vaccine doses from the same manufacturer should be administered whenever recommended. In the following circumstances, an age-appropriate COVID-19 vaccine from a different manufacturer may be administered:

- Same vaccine not available at the vaccination site at the time of the clinic visit
- Previous dose unknown
- Person would otherwise not receive a recommended vaccine dose
- Person starts but unable to complete a vaccination series with the same COVID-19 vaccine due to a contraindication